

Mr Malcolm Cameron FRCS(OMFS) FRCS(Eng) FDSRCS(Eng)
Consultant Head & Neck Maxillofacial Surgeon

The Nuffield Health Cambridge Hospital
4 Trumpington Road
Cambridge
CB2 8AF

All enquires & correspondence to:
Secretary to Mr Malcolm Cameron
Spire Cambridge Lea Hospital

Tel/Fax: 01223 266928
mmp2@themedman

Spire Cambridge Lea Hospital
30 New Road
Impington
Cambridge, CB24 9EL

Appointments: 01223 266990

REMOVAL OF IMPACTED WISDOM TEETH

This leaflet has been designed to improve your understanding of your forthcoming treatment and contains answers to many of the commonly asked questions. If you have any other questions that the leaflet does not answer or would like further explanation please ask.

The problem

Third molars, better known as wisdom teeth, are normally the last teeth to erupt into the mouth and typically do so between the ages of 18 and 24. They are the eighth tooth back in each corner of the mouth so also known as “8’s”. In modern populations, there is often not enough room to allow the wisdom teeth to come through properly and when this happens, they are said to be “impacted”. Consequences of this include recurrent infections, food packing and other problems.

Why do I need treatment?

Impacted wisdom teeth can cause a number of problems that mean that the tooth or teeth are best removed. The most common problems include:

- Recurring infection around the crown of the tooth. This is known as pericoronitis and leads to pain and swelling.
- Decay in the wisdom tooth or in the tooth in front.
- Food packing.
- Cysts can form around the wisdom tooth if it does not erupt into the mouth properly. A cyst occurs when fluid fills the sack that normally surrounds a developing wisdom tooth.

What does the treatment involve?

Because the wisdom tooth has not fully erupted it is often necessary to make a cut in the gum adjacent to the tooth. Sometimes it is also necessary to remove some bone surrounding the crown of the wisdom tooth. The wisdom tooth may need to be cut into

pieces to remove it. Once the wisdom tooth has been removed the gum is put back into place with stitches. These stitches are dissolvable and take around two weeks to disappear.

What type of anaesthetic is used?

Several options are available and depend on how difficult the wisdom tooth is to remove:

- **Local anaesthetic** - this is an injection into gum surrounding the wisdom tooth, similar to that you may have had at your dentist for a filling. The injection takes a couple of minutes to work and numb the area and means that you will feel no pain while the wisdom tooth is removed. This is the best option for wisdom teeth that are straight forward to remove.
- **Local anaesthetic and intravenous sedation** – in addition to a local anaesthetic injection you can be given an injection into your arm. This makes you feel relaxed and less aware of the procedure.
- **General anaesthetic** – it is usually possible to remove wisdom teeth under a “day case” general anaesthetic, ie you are put to sleep completely and will be able to go home on the same day.

How long does it take to remove a wisdom tooth?

This is a variable. Some wisdom teeth may take only a few minutes to remove. More difficult wisdom teeth that need to be cut into pieces to remove can take around 20 minutes or more to extract.

Is there much pain or swelling after the removal of wisdom teeth?

It is likely that there will be some discomfort and swelling both on the inside and outside of your mouth after surgery. This is usually worse for the first three days and may take up to two weeks before all the pain goes. You may also find that your jaw is stiff and you may need to eat a soft diet for a week or so. If it is likely to be sore, you will be given painkillers. It may also be necessary for you to have a course of antibiotics after the extraction. There may be some bruising of the skin of your face that can take up to a fortnight to fade away.

Is there anything else I need to do after the extractions?

It is important to keep the extraction sites as clean as possible after the surgery. It may be difficult to clean your teeth around the sites of the extraction because it is sore and if this is the case it is best to keep the area free from food debris by gently rinsing with a

mouthwash or warm salt water (dissolve a flat teaspoon of kitchen salt in a cup of warm water) commencing on the day after surgery.

Do I need to take any time off work?

Usually, it will be necessary to take a few days off work and avoid strenuous exercise for this time. You should not drive for 24 hours after intravenous sedation and for 48 hours after a general anaesthetic.

What are the possible problems?

- Although there may be a little bleeding at the time of the extraction this usually stops very quickly and is unlikely to be a problem if the wound is stitched. Should the area bleed again when you get home this can usually be stopped by applying pressure over the area for at least 15 minutes with a rolled up handkerchief or swab. If the bleeding does not stop, please get in contact.
- Infection is uncommon particularly if antibiotics are used.
- There are two nerves that lie very close to the roots of the lower wisdom teeth. One of these nerves supplies feeling to your lower lip, chin and lower teeth. The other supplies feeling to your tongue and helps with taste. Sometimes these nerves may be bruised when a wisdom tooth is taken out. This can cause tingling or numbness in your lip, chin or tongue, and more rarely cause altered taste. About one in 10 people will have some tingling or numbness that can last several weeks. Less than one in 100 people will have problems that last more than a year. These risks may be higher if your tooth is in a difficult position. The surgeon will tell you if you are considered to be at an increased risk.

For further advice and information please see the web pages of The British Association of Oral & Maxillofacial Surgeons www.baoms.org.uk